

## Kansas Immunization Requirement School Nurse Tool – 2021-22 School Year [K.C.I.](#)

Vaccine doses administered **up to 4 days** before the minimum interval of age can be **counted as valid** with the **exception of two live vaccines which require 28 days between doses**. A child must be at least six weeks of age prior to receiving vaccines (only exception is Hepatitis B). Each bulleted criterion must be met for stated number of doses to meet requirement. Month = 28 days for intervals < 4 months. Month = calendar month for intervals ≥ 4 months.

Doses	
# required	<b>DTaP/DT</b> - Limit 6 doses for DTaP (5 <sup>th</sup> dose is not necessary if 4 <sup>th</sup> dose is ≥ 4 <sup>th</sup> birthday and ≥ 6 months after 3 <sup>rd</sup> dose.)
4 or 5 doses	<ul style="list-style-type: none"> <li>First dose at least 6 weeks old</li> <li>At least 4 weeks between doses except -                             <ul style="list-style-type: none"> <li>At least <b>6 months* between #3 &amp; #4 and #4 &amp; #5</b> (*four months acceptable between dose 3 &amp; 4, but not recommended)</li> <li>One dose on or after 4<sup>th</sup> birthday (<i>Tdap given to a child younger than age 7 years as either dose 1, 2, or 3, is NOT valid. Repeat with DTaP as soon as feasible. Tdap given as either dose 4 or 5 is valid.</i>) <b>Note:</b> DTaP not recommended for children over age 7 due to local irritation.</li> </ul> </li> </ul>
<b>Td/Tdap – delayed/makeup schedule – child ≥7 years only</b>	
3 doses	<ul style="list-style-type: none"> <li>First dose should be Tdap for the delayed/makeup schedule (if child is ≥ 7 years)</li> <li>At least 4 weeks between doses 1 &amp; 2 <b>and</b> at least 6 months between doses 2 &amp; 3</li> </ul>
<b>Note:</b> DTaP given from ages 10-18 may count as Tdap	
<b>Note:</b> A single dose of Tdap can be counted for an incomplete primary DTaP series if child is ≥7 years and lacking only 1 dose.	
<b>Note:</b> If child age ≥7 completing the Td/Tdap series received any doses of DTaP/DT prior to age 12 months, ≥ four doses are required.	
<b>Tdap</b>	
1 dose	Proof of 1 Tdap booster required for Grades 7 – 12 <i>Children aged 7-9 receiving Tdap should get routine Tdap at ages 11-12, but children aged 10 receiving Tdap do not need routine dose at ages 11-12.</i>
<b>MMR</b>	
2 doses	<ul style="list-style-type: none"> <li>1<sup>st</sup> dose on or after first birthday</li> <li>At least 28 days between doses</li> </ul>
<b>Varicella</b>	
1 dose	<ul style="list-style-type: none"> <li>Required on or after first birthday</li> </ul>
2 doses	<ul style="list-style-type: none"> <li>Required for Grades K – 12</li> <li><b>If younger than age 13</b>, 3 months between doses preferred (28 days accepted). <b>If 13 or older</b>, ≥28 days between doses.</li> </ul>
Disease history accepted with physician or health department signature <b>or</b> lab verification of positive titers.	

Doses	
# required	<b>Polio</b> IPV not used prior to 1997. <b>OPV given on or after April 1, 2016 should not be counted.</b>
4 doses	Any combination of OPV/IPV or all OPV or all IPV <ul style="list-style-type: none"> <li>First dose at least 6 weeks old</li> <li>At least 4 weeks between doses (<b>Grades K - 10 are required to have six months between doses 3 and 4.</b>)</li> <li><b>Grades K - 10, one dose required on or after 4<sup>th</sup> birthday regardless of number of prior doses.</b></li> </ul>
3 doses	All 3 OPV or all 3 IPV <ul style="list-style-type: none"> <li>First dose at least 6 weeks old</li> <li>At least 4 weeks between doses (<b>Grades K – 10 must have six months between doses 2 and 3.</b>)</li> <li>One dose on or after 4<sup>th</sup> birthday</li> <li>Older children needing to complete the series require 6 months between doses 2 and 3</li> </ul>
<b>Hepatitis B</b>	
<b>Note:</b> Four doses of Hep B can be given when a birth dose is given.	
3 doses	<ul style="list-style-type: none"> <li>1<sup>st</sup> dose can be given as early as day of birth</li> <li>At least 4 weeks between dose 1 &amp; 2</li> <li>At least 8 weeks between dose 2 &amp; 3</li> <li>Must be a total of 16 weeks between dose 1 &amp; 3 <b>and</b> ≥ 6 months of age for dose 3</li> </ul>
2 doses	<ul style="list-style-type: none"> <li>Recombivax 1 cc give between 11 to 15 years of age <b>and</b></li> <li>Dose 1 and 2 separated by 4 to 6 months</li> </ul>
<b>Gardasil (HPV) (Not required – strongly recommended)</b>	
2 doses	<ul style="list-style-type: none"> <li>Minimum age – 9 yrs. Prefer age 11 or older</li> <li>If &lt; 15 when receiving first dose only 2 doses are needed (6 months between doses)</li> </ul>
3 doses	<ul style="list-style-type: none"> <li>If &gt;15 for first dose require minimum 4 weeks between dose 1 and 2 and 6 months between dose 1 and 3</li> </ul>

<b>Doses</b>	<b>Hepatitis A</b> (Required for Early Childhood through <b>Grade 5</b> )	<b>Doses</b>	<b>Meningococcal A,C,W,Y</b> (Required for Grades <b>7-9</b> & 11-12)
2 doses	<ul style="list-style-type: none"> <li>1<sup>st</sup> dose on or after first birthday</li> </ul>	2 doses	1 <sup>st</sup> dose required at Grade 7 ( 2 <sup>nd</sup> dose due Grade 11)
	<ul style="list-style-type: none"> <li>Must be 6 months between doses</li> </ul>		2 <sup>nd</sup> dose required at Grade 11 (age 16-18) (see 1-dose age exception below)
	Recommended for <i>all</i> children age 1 year and up.	1 dose	<b>If 1<sup>st</sup> dose administered at ≥ 16, does not need 2<sup>nd</sup> dose.</b>
	<b>Haemophilus influenzae type B* (Hib)</b> (Required for Early Childhood). If Hib unknown, follow Act HIB schedule.		<b>Pneumococcal Conjugate**</b> (Required for Early Childhood)
4 doses	The usual schedule for <b>ActHIB is 4 doses</b> with first three doses given at 2, 4 and 6 months and the 4 <sup>th</sup> dose given at 12 to 15 months of age. Spacing requirements are as follows: <ul style="list-style-type: none"> <li>At least 4 weeks between first 3 doses</li> <li>8 weeks between dose 3 and 4</li> <li>4<sup>th</sup> dose on or after 1<sup>st</sup> birthday</li> </ul>	4 doses	Usual schedule <b>is 4 doses</b> with first three doses given at 2, 4 and 6 months and the 4 <sup>th</sup> dose given at 12 to 15 months of age. Spacing requirements are as follows: <ul style="list-style-type: none"> <li>At least 4 weeks between first 3 doses</li> <li>8 weeks between dose 3 and 4</li> <li>4<sup>th</sup> dose on or after 1<sup>st</sup> birthday</li> </ul>
3 doses	The usual schedule for <b>PedvaxHIB is 3 doses</b> with first two doses given at 2 and 4 months (minimal interval of 4 weeks) and the 3 <sup>rd</sup> dose give at 12 to 15 months of age (minimal interval 8 weeks from last dose).	3 doses	<ul style="list-style-type: none"> <li>At least 4 weeks between first 2 doses</li> <li>8 weeks between dose 2 and 3</li> <li>3<sup>rd</sup> dose must be on or after 1<sup>st</sup> birthday (completes series in healthy child)</li> </ul>
3 doses	<b>If age 7-11 months when receiving first dose there must be:</b> <ul style="list-style-type: none"> <li>Four weeks between first 2 doses</li> <li>8 weeks between dose 2 and 3</li> <li>3<sup>rd</sup> dose on or after 1<sup>st</sup> birthday</li> </ul>	2 doses	<b>If age 12 to 23 months when receiving first dose there must be:</b> <ul style="list-style-type: none"> <li>At least 8 weeks between doses</li> </ul>
2 doses	<b>If age 12 to 14 months when receiving first dose there must be:</b> <ul style="list-style-type: none"> <li>1<sup>st</sup> dose given between 12 up to 14 months of age</li> <li>8 weeks between doses</li> </ul>	1 dose	<b>If age 24 months up to 59 months when receiving first dose</b> only one dose required. <b>NOTE:</b> If one dose has been received <b>after age 24 months</b> the child is <b>done</b> if at least 8 weeks between the most recent prior dose (if any)
1 dose	<b>If age 15 months up to 59 months when receiving first dose</b> only one dose required. <b>NOTE:</b> If one dose has been received <b>after age 15 months</b> the child is <b>done</b> if at least 8 weeks between the most recent prior dose (if any)	2 doses	<b>Note:</b> Unvaccinated children 24-71 months of age <u>with</u> underlying medical conditions should receive 2 doses separated by at least 8 weeks.
	*In general, Hib vaccination is not routinely recommended for persons older than 59 months of age.		**Pneumococcal Conjugate is not routinely recommended for persons older than 59 months of age.

Refer to CDC Catch Up Schedule <http://www.cdc.gov/vaccines/schedules/hcp/imz/catchup.html>

Table Tool created by Olathe Public Schools (USD 233) in consultation with the Kansas School Nurses Organization Rev. 4/21